



Advances in Predictive Modeling: The Role of Artificial Intelligence in Monitoring Blood Lactate Levels Post-Cardiac Surgery

Roman Khan¹, Hira Zainab², Arbaz Haider Khan³, Hafiz Khawar Hussain⁴

¹Lewis University Chicago

²Department of Information Technology American National University

³University of Punjab

⁴DePaul University Chicago, Illinois

¹romankhan@lewisu.edu, ²hira.zainab72@gmail.com, ³arbazhaiderkhan15@gmail.com, ⁴Hhussa14@depaul.edu



Corresponding Author

Hafiz Khawar Hussain

Hhussa14@depaul.edu

Article History:

Submitted: 14-11-2024

Accepted: 14-11-2024

Published: 14-11-2024

Keywords: AI, lactate tracking, cardiac procedures, intensive care, prognostic analysis, effectiveness, patients' experiences, Intensive Care Unit, live data, big data, alerting, critical care, advanced technologies, ethical issues, possible overtones, overstimulation, medicines for individuals

Brilliance: Research of

Artificial Intelligence is licensed

under a Creative Commons

Attribution-Noncommercial 4.0

International (CC BY-NC 4.0).

ABSTRACT

Total blood lactate levels monitoring through the use of Artificial Intelligence in individuals that have undergone cardio surgeries is a milestone in critical care because it indicates metabolic problems earlier than traditional approaches. Lactate levels have to be significantly raised in order they may indicate complications like tissue hypoxia, sepsis or organ dysfunction. The previous method of monitoring lactate entails conducting tests after a few hours or days and can be very unresponsive; in the application of AI models, the algorithm scans through data acquired from patient monitoring systems to predict and advance notice the clinicians on the trends in lactate levels. This review outlines the basic mechanisms, algorithms, and features required to build an AI-based lactate predictor and the multiple physiologic signals such as heart rate, oxygen saturation, and blood pressure into the support vector regression model. Illustrative cases show that AI can facilitate more effective clinical decision-making to increase ICU patient safety and decrease such hospital stays. While AI based lactate tracking is something that has been bandied about in the research literature for some time, there are real questions as to how this is implemented in existing hospitals, how one minimizes the negative impacts of alarm fatigue, and how the results are persistent across population groups. Ethical and legal necessities concerning patient's data confidentiality, security, and further reporting also play the vital role of its clinical endorsement. Other directions for future work are more flexible and multiple modality models that include additional data and require learning from new patient data.

INTRODUCTION

Introducing an article on recent developments in predictive modeling with considerable emphasis on post-cardiac surgical blood lactate monitoring using artificial intelligence (AI), raises the awareness of why lactate monitoring is so important in the clinical management of patients after complex surgeries. Lactate is better known, and typically utilized as the primary indicator when tracking critically ill patients and can lead to tissue hypoxia as well as metabolic acidosis, sepsis or organ failure, especially in the post-surgical care of a cardiac surgery patient. Lactate measurement allows clinicians to detect the onset of complications, change management as necessary, and achieve the best results [1]. However, conventional methods for lactate assessment and forecasting are slow, invasive, and usually do not provide enough accuracy for estimating the likelihood of complications before they occur. This has given rise to a higher emphasis on the predictive modeling area, the focus on which is driven by the AI solutions.

Blood lactate monitoring should be conducted in the early period after cardiac surgery because many of such surgeries can be high-risk associated with long anesthesia time and potential ischemic episodes. The level of lactate in the blood helps to figure out whether tissues are well oxygenated and whether cells experience anaerobic metabolism [2]. When cells shift to anaerobic pathways for energy generation, lactate is one of the by-products in this process, and identifying high level of the substance in the blood stream determine that a patient is suffering from hypoxia, this is a severe state that requires the attention of health practitioners. This may be especially true for lactate monitoring post-surgery, because while its values are being closely monitored they may be in ICU where numerous other physiological parameters abound [3]. Lactate has been traditionally assessed by drawing a blood sample from the patient which is inconvenient, costly and time consuming, and the sampling frequency was limited. This approach may not detect changes in lactate levels that may





point to the onset of new clinical problems. Hence, the call for enhanced models of more dynamic lactate changes that would help predict these changes in advance thus enabling care givers to prevent such changes from occurring.

However, with lactate monitoring considered such a critical factor, there are several difficulties inherent in traditional methods. First of all, that is, due to the actual periodicity of blood sampling, it is possible to measure lactate levels only occasionally. This may result to the practitioner taking longer time to note increased levels of lactate; hence, the missed opportunity to intervene timely. Moreover, within patient variables, comorbid illness and concomitant medications make definition of a normal value for this population problematic [4]. Evaluation of lactate patterns needs clinical experience and can be next to impossible due to human intervention and mistake particularly in a busy ICU where several crucial markers have to be observed. Another limitation of the approach that traditional monitoring offers is that it does not offer a predictive vantage. Conventional procedures are involved in monitoring of lactate levels in the body without giving estimates of the fluctuations which may occur in near future. The importance is best summarized by the fact that when it comes to adverse trends, early detection is as crucial as early intervention is helpful and it is for this reason that tracking tools – not only for lactate levels – but also for the likelihood of a rise in lactate levels, are invaluable [5].


Over the last few years, the advances made in the area of predictive modeling especially through AI technologies have brought exciting innovations to the field of health surveillance. AI system can well analyze enormous patient records, detect the repetitive data pattern and come up with the forecasts, which are unattainable by conventional approach solely. This means, that, for instance, when considering the post-cardiac surgery lactate series, AI-based models can leverage numeric values, such as heart rate, blood pressure, oxygen saturation, or even patient's demographics to predict changes in lactate levels ahead of their occurrence. Used in early- warning system, machine learning algorithms can be trained to identify even tough obscure patterns and correlations which human mind cannot decipher [6]. Artificial intelligence based lactate prediction models could influence post-operative care by presenting accurate, 'in the moment', lactate prediction for use in these decisions. Such models must be supervised with data which they are taught with being historical patient data in order that they may increase their reliability and specifically generate accurate individual patient predictions. The use of AI in managing lactate levels is best suited in the ICU, because predictive tools would help in managing patient flow, organizing patients' treatment and care hence can replace some of the burdens of healthcare workers.

The resulting change to a more habitual concentration on predicting lactate patterns means cardiac surgery patients can benefit greatly. By identifying that the lactate level is rising early one can actually change certain medications, fluids or oxygen delivery to bring the condition of the patient back into normal range. Moreover, AI-based lactate monitoring systems offer opportunities to create scenarios for continuous and non-invasive monitoring systems, excluding potential dangers for patients linked to invasions of blood samples [7]. With the further development of AI technologies, integrating them into the functional processes of an ICU and a surgical recovery stage will be even more uninhibited allowing to include the predictive lactate monitoring into the standard of post-Cardiac surgery management. To a summary, the use of AI can bring statics beyond those currently available in the practices of lactate monitoring, providing timely, accurate, and proactive patient data post-Cardiac surgery. Subsequent divisions of this article will discuss future sections and examine the prevailing technologies, algorithms, and clinical uses of AI for lactate prediction in correlation to its constructive and constraining factors on advanced cardiology.

PREDICTIVE MODELING IN CARDIAC SURGERY

Clinical decision support has thus been introduced as a new tool in which predictive modeling is named as a distinct avenue in healthcare, which gives clinicians the possibility to forecast the outcome and act accordingly. Thus, the value of predictive models is high in such surgical procedures as cardiac surgery where patients' risk is high and postoperative course may be lengthy and pose many challenges [8]. These models can be really useful as by analyzing data which was collected earlier and looking for patterns in patients' physiology and their response towards certain treatments, the specialists will be able to foresee certain complications, manage the interventions and, therefore, will be able to affect the overall result. Section 2 and its subdivisions provide an overview about the historical development of PM as well as the application of this technique in the context of cardiac surgical patient management and the clinical factors that are important for the prediction of blood lactate concentrations after surgery in particular [9].

Emerging trends of Predictive Models in Medicine: Another review article Mosteller pointed out that predictive modeling in medicine has progressed notably in recent decades, primarily due to improvements in statistical methodologies and in data analysis science. The initial frameworks depended on logistic regression and other analytical tools in order to make estimations with the help of a small number of characteristics. With the rising of big data and application of ML techniques, the techniques has expanded the component types by including not only the gene markers but also the real time physiological data for much fine-grained prediction models. Neural networks, support vector machines and decision trees and other advanced technology has helped in developing predictive models that can work on non-linearity relationships among different elements [10]. For example, predicting postoperative course after cardiac surgery is dependent on a number of variables which will include but not be limited to the patient's age, previous medical history, intraoperative variables and postoperative variables like the patient blood pressure and oxygen saturation levels. Using these variables, the ML algorithms can assign weights to them in dependence with their correlation and then headers; can give inferences that might show the potential of a specific patient to have complications or early





interventions.

The Urgency of Using the Predictive Modeling Systems in Cardiac Surgical Practice: They claim that morbidity and mortality of cardiac surgery are significant driven by the complex process of operating on the heart and vascular system. Operations such as CABG, valve replacements, or valve repairs for example, include durations of controlled ischemia, heart & lung bypass, and anesthesia all of which challenge the metabolic economies of the body. Cardiac arrhythmias, infection, hemorrhage, and organ dysfunction are not rare events in the postoperative period and need constant supervision. Blood lactate concentrations are one of the variables which is used to measure metabolic pressure and tissue oxygenation [11]. However, the traditional approaches in monitoring are reactive in nature, revealing increased lactate concentrations once the situation worsens, constraining the means for clinicians to prevent compounding of those.

Predictive modelling has an answer to this challenge by highlighting the number of initial indicators that work as predictors, showing which of the patients is more likely to encounter problems, so that intervention is made early enough and passed through the most effective personalized treatment plan. Further, it supports risk categorization of patients, to enhance the future utilization and efficiency of actual ICU, by directing it resources to those most deserving. It is certainly capable of bettering the overall outcome of patient care but can also address issues of productivity in the limited environment of a hospital [12]. Using severe sepsis scoring systems to predict the level of lactate in the post-operative patient is analysed in terms of the following key clinical parameters; most important, to predict blood lactate levels after cardiac surgery, it is necessary to collect many clinical factors that impact on the metabolism and oxygenation of tissues. Some of the vital patient parameters are the blood pressure, pulse rate, oxygen saturation which help index the patient's circulatory and respiratory condition respectively. Specifically, hypotension and hypoxaemia are important factors underlying tissue hypoxia and subsequent anaerobic processes with increased production of lactate [13].

Other potential parameters are variables related to the surgical procedures like time of surgery, the time on cardiopulmonary bypass and blood loss. For example, duration of surgery and bypass time have certain direct correlation with metabolic complications observed in patient's condition. In addition, blood transfusion, medication and fluid balance administered before, during and after surgical procedure can affect the lactate levels. Surprisingly, even the patient-specific variables including the patient's age, BMI, past medical history (diabetes/chronic renal disease), and basic metabolic profile were found relevant in determining postoperative lactate kinetics. Lately, the "input variables" have included laboratory test results, e.g., hemoglobin, white cell count, serum electrolytes as well as the changes thereof within the past short period. Micro process techniques, such as ABG and near-infrared spectroscopy are in contrast comparatively more dynamic in their approach to identifying the metabolic condition of the patient [14]. When integrated with predictive analytics, such technologies will help in offering real-time lactate level estimates for a patient to facilitate a much more effective clinical intervention.

Some of the key challenges involved in work on integrating predictive models into care of patients undergoing cardiac surgery include not only building good-enough models, but also incorporating these models in the workflow in a manner that adds value to rather than encumbering the work of point-of-care clinicians. Structured models for lactate tracking for instance can be incorporated into EHRs and/or ICU dashboard where it will include an alert when observed lactate probabilities are potentially hazardous [15]. These alerts give specific, targeted information so that clinicians can change their approach at the next step, for example, to give more oxygen, to take steps to better deal with fluid status, or to modify drugs to enhance tissue perfusion. As such, predictive models appear to have the potential to revolutionize's post-operative care in cardiac surgery by improving clinicians' ability to predict unfavorable outcomes. Still, since the use of these tools involves problematic validation issues and direct interaction with healthcare providers, their potential positive effect on reported patient outcomes and healthcare production cannot be questioned [16].

APPLICATIONS OF ARTIFICIAL INTELLIGENCE IN HEALTHCARE AND MACHINE LEARNING

Artificial intelligence (AI) and machine learning (ML) having emerged as disruptive technologies, continue to gain importance in transforming the application of data for prognosis with regards to health. In critical care and postoperative care, AI and ML find use in tracking patients' conditions and identifying factors that put patients at risk for developed complications. Their application after cardiac surgery to establish blood lactate concentration is an illustration of the manner in which such tools guide carry invaluable understanding of the patient's physiology and help professionals to foretell adverse events. In this section, the author provides a taxonomy of the AI and ML approaches that can be used in health care, especially regarding post-operative observations of the patient and reviews the benefits and drawbacks of the integration of artificial intelligence-driven solutions in medical practice [17]. Machine learning and artificial neural network can be defined as the use of computational methods to make machines able to process data, to recognize a pattern or some dependencies, to perform prediction or to make decisions with the help of received data. Artificial Intelligence was defined in this research study as a branch of computer science involving the use of statistical techniques to learn from data to recognize patterns without being programmed. The basic categories of machine learning are also valuable in the healthcare industry: supervised, unsupervised, reinforcement, and deep learning. Et al., showed that each approach is helpful in different cases depending on the type of data present and problem being solved in clinical setting.

The algorithms such as supervised learning is trained on a labelled dataset and often performs very well if there is very little noise. These models can learn the correlation of, for example, heart rate, blood pressure or SpO₂ and lactate or potential





complications [18]. Structural learning, in contrast, is applied to identify hidden patterns or clustering throughout the datasets, which may be useful when the identification of risk-profiles of subgroups of patients exists. The second a bit more complex is the neural networks based technique known as deep learning that is perfect for processing of large and complex data of various origin including imaging or multi-parameter ICU monitoring system.


Reinforcement learning is also showing its applicability to healthcare, particularly to improve the treatment plan. On this approach, algorithms gain feedback from the decisions that they make, and change with the results. Reinforcement learning seems relevant in the complex and ever-evolving context of the ICU, with the many different conditions to constantly learn from clinical data and with ongoing feedback. The integration of AI and ML in health care but mainly in intensive care as well as postoperative care unit has the following advantages. Perhaps the most important is the capacity of artificial intelligence models to identify the correlation between different parameters of a patient's state which cannot be noticed by a doctor during several visits, as well as to analyze the data of countless patients in real time [19]. It is particularly indispensable in tracking multiple physiological parameters that are influenced by multiple factors in a patient's organism such as altering lactate levels in patient's organism.

AI models when providing predictive analytics can provide information quicker and with higher accuracy compared to conventional approaches and assist in early stratification to enhance patients' prognosis. For instance, if an AI model indicates lactate levels are going to increase then healthcare providers can manage for anticipated issues that come with increased lactate levels such as tissue hypoxia or sepsis before they occur. Moreover, the AI-based systems help the opposite of overwhelming the human healthcare staff as they take some shares of the data analysis and provide human workers with more effective tools for decision-making [20]. A benefit is that through AI technology, patient healthcare can be given individually. The utilization of patient associated clinical data fulfills the needs of AI models where specific patient diagnosis and prognosis needs individualized predictions of towards best treatment approaches that considers case factors like age, historical health records and family genetics. Patients undergoing cardiac surgery have different levels of comorbidity, and individual risk assessment would significantly improve post-operative surveillance. It is therefore important to discuss the limitations of AI in the clinical scenario and continued challenges that remain largely unresolved [21].

In the following section, we outline various challenges that healthcare and their stakeholders stand to face after integrating AI technology. As one of the biggest challenges, it is still questionable what kind of data is delivered and how this data varies. Clinical data are frequently siloed, partial, and heterogeneous across care settings which raises the question of how AI models perform accurately and can generalize beyond them. For example, Data collected from EHRs might be erroneous or not adhere to a common guideline, leading to questions marks on the validity of the predictions made in the triage of patients in the intensive care unit. They include the raised privacy concerns associated with the 'black box' characterization of some of the most contemporary AI models – namely, deeper learning algorithms [22]. Many of these models can provide solutions with great precision but are typically insufficient in justifying the steps they took to come up with those answers; this makes it hard for clinicians to even have faith in their results or even base their decisions on them. This lack of interpretability particularly becomes a hinderance in delicate atmosphere such as after cardiac surgery, a clinician cannot be dragged in making a decision based on such analysis and probabilities.

However, ethical and regulatory considerations are still questions that need to be solved on the way to using artificial intelligence in clinics. Privacy issues cannot be overlooked since, for most of the AI models, patients' data is needed to achieve high predictive accuracy. These and many other issues indicate the need for AI systems to be developed in full compliance with regulatory acts like HIPAA [23]. AI steps into the healthcare system to support clinical decisions, there is another question that needs to be answered – who is culpable for the deleterious consequences when such recommendation has been provided by an AI source? Last but not least, the deployment of AI tools is expensive in terms of technology as well as human resource development. We found that using AI in hospitals and clinics require development of infrastructure that is capable of handling real-time processing of data, and that healthcare providers need to be taught on how to integrate AI systems effectively into their working environment. By following the machine decision but not the analysis, there is a possibility that clinicians will always believe in the machine or always disregard them because they do not understand them or do not trust them.

Big Data, AI, and ML are getting the power to enhance healthcare, especially when it measures constant attention and timely examination is critical, such as intense care units. This paper demonstrated that if integrated in patient care post-operative monitoring of patients who underwent cardiac surgery could become more proactive, precise and responsive since AI offers data-driven predictions that support individualised care [24]. However, for these models to reach their fourth dimension, issues regarding the data, model, its explanation, the ethical use, and infrastructural requirements have to be solved. The proper integration of these technologies into healthcare urgently requires standardization of AI technologies and the determination of best practices for their use.





BREAKDOWN OF TYPES OF DOCUMENTATION ISSUES IN HIGH-SEVERITY INPATIENT SURGICAL TREATMENT CLAIMS





Figure: 1 showing Breakdown of types of documentation issues in high-severity inpatient surgical treatment claims in percentage

APPROACHES FOR AI MODELS FORECASTING LACTATE LEVEL

The employment of AI and ML algorithms to predict the values of blood lactates post-cardiac surgery appears to be a valuable and useful idea to improve the safety and effectiveness of postoperative surveillance. Blood lactate is a very sensitive, specific indicator of patient’s metabolic status and tissue oxygenation which are particularly relevant in major surgery including cardiac surgery. Increased serum lactate in a patient is often associated with conditions like tissue hypoxia, sepsis or an organ dysfunction which should therefore be managed promptly. This section focuses on the aspect of data sources, the major AI methodologies, and feature selection on Lactate level prediction which plays a paramount role in the application of Artificial Intelligence in transforming the real-time monitoring process and the subsequent proactive patient management. More specific and general parameters have to be included in CLM in order to obtain accurate lactate prediction, in view of various aspects of physiological and clinical factors influencing the metabolic and circulatory systems. The guidelines often employed in the AI-dependent lactate estimation are abstracted from Electronic Health Records (EHRs), real-time surveillance devices, and lab values [25]. The most relevant data include the assessment





of patient's physiological state and the objective indexes such as heart rate, blood pressure, oxygen saturation, pH levels, bicarbonate, hemoglobin, while surgical data include the type of surgery, duration of the cardiopulmonary bypass, intraoperative fluid balance, and blood products transfused.

Data cleaning is the process of removing noise and converting data into a format that is ready for analysis cleaning of data can be of two types: data preprocessing and data reproduction. This step could involve dealing with missing values or extreme values which need reduction or handling, normalization that is the reduction of variability and handling of data that is inconsistent between data sets. By this, the author exhibits the importance of the quality and consistency of the data with which AI operates since the latter is negatively affected by low-quality data and noise. In the same process, the other techniques of feature engineering are also employed for the generation of new features that captures different patterns similar to trends in oxygen and or lactate levels or, the difference of lactate over some time [26]. These engineered features then can feed some useful information to the AI model that in return help in the construction of the best predictive model. Several forms of ML & AI approaches can be used to forecast lactate levels; each has its advantages and disadvantages. Some of the commonly used algorithms are Linear, Logistic Regression etc which are conventional and serve little in big data and yet offer simple output. They may be not sufficient to measure the non-linear interactions among multiple variables impacting lactate kinetics [27].

For more detailed and curvilinear data and working with discrete data, such as decision trees, support vector machines (SVM), and some types of ensemble methods (random forests and gradient boosting, for example). These models work well with more datasets and more predictors making it possible to understand how the different variables influence the lactate level change. For instance, random forests not only can identify combined effects among variables, including oxygen saturation, fluid management, and blood pressure on lactate levels, but also assign the rankings of each variable's importance for the prediction. Neural networks and deep learning algorithms are most useful for more complex kinds of monitoring where trend data is available for ongoing review such as in Intensive Care Units (ICUs) [28]. The type of ANN especially suited for predicting lactate levels is the recurrent neural networks and their variants including LSTM because they are trained to handle sequences of data over time in order to capture dependency on time from a patient's monitored vitals and laboratory results. For example, an LSTM-based model can take time dependencies in each of heart rate, respiratory rate, and oxygen level to discover unknown patterns that show an upcoming increase in lactate.

Thus, one of the major problems when constructing the lactate predictive model is the choice of features – variables that best describe the potential of a patient to have an elevated lactate levels. Feature selection it is a process where variables relevant to the data set are singled out in a bid to remove noise to make the model more accurate [29]. Some of the most basic approaches include obtaining feature subsets that are highly correlated with lactate levels and more sophisticated approaches which are recurrent feature elimination and others. It could be stated that the most important features for lactate prediction might include both vital signs and laboratory results and surgical characteristics. For example, mean arterial pressure, arterial blood oxygen, haemoglobin concentration, and blood pH all reflect tissue oxygen delivery, and the metabolic rate, which are crucial determinants of lactate concentrations. Further, some variables, including fluctuation of key parameters over a given time interval and trends in these changes, can signal about physiological stress before elevation of lactate, for example, fluctuations in the heart rate. Also, the AI models can rank these features according to their contribution level; this information is precious on a clinical aspect [30]. For instance a model might show that the most important variables that predict lactate rise in the hours following surgery are oxygen saturation and blood pressures implying that these should be strictly monitored. They also present a use in improving the model itself, in that they inform the practitioner which variables are the most influential for accurate prediction, and could soon lead to the development of new models that are simpler and more easily integrated into clinical practice.

Once an AI model is trained and validated it can be the used in the real time surveillance systems in the ICU or other post-surgical care settings. The model is constantly collecting data from patient monitors and laboratory systems and continuously provides ongoing prognosis of lactate level increase [31]. In real-time applications, the AI model offers clinicians a decision support system that gives alerts or risk scores that trigger appropriate action that may include changing oxygen rate, fluid balance or medications aimed at stabilizing the patient. The use of MMP and SP for accommodation in the clinical context particular to the ICU has the following benefits. This makes it facilitate a more predictive entry of patient care where frequent blood samples are not needed, rather constant blood glucose testing. If clinicians receive audible alert of possible rise in lactate, they attend to the patients early enough to attend to the cause, and possibly reversing any more rise in lactate thus enhancing patient outcomes and possibly the short hospital stays in the ICU. Nevertheless, these AI-driven models can be effectively used for more efficient ICU internal resource distribution as well as alarming of dangerous prognosis patients and their closer monitoring, which will contribute to the general increase in productivity of the ICU [32].

The use of machine learning algorithms for the prediction of lactate levels entails a new approach to the clinical practice of the post-cardiac surgery monitoring. Whereas the traditional analysis of clinical databases often provides a single picture of a patient's condition, these models can use various databases and better algorithms to make more precise and instant predictions for clinical decision-making. Both the feature selection strategy and the real-time data integration process are critical to success of these models and should be capable of providing good robust physiological signal coverage's that fit well into the current clinical and operating environment of the clinical setting [33]. Thus as the





prediction models continue to be developed the use of AI in the patient care systems can only prove to be beneficial as it will improve the chances of favorable outcomes for the patients as well as tripling the efficiency of treatment in intensive care units.

CLINICALLY INCLUDING APPLICATIONS AND CASES OF AI IN LACTATE MONITORING

AI and ML in lactate monitoring have been developed to improve lactate measurement in clinical contexts, especially for patients who have been operated using risky cardiac surgery. Use cases based on clinical cases show the benefits of AI models for better patient outcomes, early recognition of potential complications, individual approaches to management and prevention. AI-based lactate monitoring has advanced from academic concepts to application in various hospitals and even Intensive Care Units (ICUs) through utilization of advanced predictive models in post-surgical care [34]. The real-world applications, examples, and analyzed outcomes of utilising AI in lactate monitoring are reviewed in this section.

Real Operating Use of AI in Lactate Control: AI has showed utility in the monitoring and early estimation of changes in lactate concentrations in ICU patients whose condition may change quickly. The main existing approaches for lactate monitoring are solely based on blood sampling and subsequent laboratory examinations, which can be invasive, time-consuming and which provide samples with an interval that is not sufficient for immediate actions when necessary. On the other hand, Artificial Intelligence based monitoring models use data from the patient's monitoring devices like heart rate, blood pressure, oxygen saturation, respiratory rate etc., to forecast the changes in Lactate level well in advance and would inform the clinicians about it. This change from reactive to proactive monitoring allows this team to get involved when lactate levels are not all that elevated, which might prevent poor results [35]. For instance, machines learning algorithms in the ICU monitor will be capable of analyzing the physiological data, which is processed consistently in order to find early signs of metabolic stress. Where the model produces an alert suggesting that lactate levels are going to rise, then clinicians can be informed to check up the status of the patient in order to fine tune the treatment plan; for instance, the oxygen delivery may be increased, the fluids may be closely monitored, the medicines may be given. The AI model therefore acts as a decision aiding tool in ensuring Clinicians keep off from deranging their patient's metabolism and thus possibly shorten their duration of stay in the ICU [36].

Artificial Intelligence for tracking Lactate Post Cardiac Surgical Procedures: Another case was conducted at a prestigious site of specialization in cardiac care, where the use of AI based lactate prediction models were used to track , patients after cardiac surgery. It was proposed that the described approach would help determine if predictive modeling is capable of predicting the rise in lactate levels and facilitate interventions before conventional methods can do so. This cohort-based AI model was derived from the information obtained from previous pediatric cardiac surgery patients in terms of their vital signs, laboratory, intra-operative and demographic data. The findings on the case study showed that, the proposed AI model had high accuracy in forecasting the elevation of lactate level several hours in advance [37]. Compared with the conventional method of monitoring, early alerts given by the model enabled clinicians not only to assist earlier the patients' needs but also modify timely the ventilating and circulating support. The active approach helped to decrease the rates of metabolic adverse effects, elevating its patients' condition and shortening their stay in the ICU. Also, clinicians had the back-up of the AI-driven model by which monitoring work was made far simpler and less cognitive and added to the efficiency of clinicians' work. This case demonstrated the use of AI solution in enhancing the post-operative care of surgical patients especially those considered at higher risks.

Large Scale Trial of Artificial Intelligence for ICU Lactate Detection: In another large scale cross sectional study, the accuracy of the six developed AI models in predicting lactate levels was evaluated in different ICU settings with different population of patients. This work employed a model trained using data obtained from more than one hospital, as such, the model was likely to perform well on a large variety of patient populations and conditions [38]. By focusing on a number of physiological markers, such as intricate heart rate variability, blood gases, and haemodynamic variables, the model was intended to detect high risk patients for pathological lactate increase, which is often related to sepsis, shock or organ dysfunction. The cross-sectional study established that the performance of the AI model was superior to conventional methods in both certainty and timeliness of identification. The predictions from the model have been embedded into ICU dashboards to show the risk scores and notify clinicians whenever the lactate levels were about to rise. The power in correctly early predicting the changes in lactate levels proved sepsis in the early stages and in most cases avoided organ dysfunction, which contributed massively to better patient's survival rates [39]. This case exemplified AI as a modular platform with value in addressing the evaluation of lactate and recommended that similar models could be implemented to improve patient care across multiple types and severities of acute illness.

Benefits of AI in Lactate Monitoring: The following main benefits were mentioned in the case studies based on the use of AI for monitoring of lactate levels: gain of better outcomes for patients, better decision making within clinics, and effectiveness of the ICU functioning. One of the biggest benefits is the move to preventative care where clinicians can then address changes in a patient's physiological state before these physiological changes result in clinical signs. As is usually the case in critical care, such early identification allows for early intervention to avert further deterioration and increase survival [40]. Different AI models negate the requirement for invasive testing by offering constant, invasive-free diploma monitoring solutions that improve patient satisfaction and security. Other advantages also highlighted in the case studies concerned the decrease of the stays in Intensive Care Units. By identifying complications before a metabolic stress





becomes worse, and lactate levels increase, the prevention of deterioration of patients facilitated by existent AI-driven lactate monitoring models, translated into shorter length of ICU stay. This means a positive impact on the health care systems since the intension of ICU overcrowding and enhanced use of its resources is addressed [41].

Real-World Implementation, Real-World Problems and Prospects: The case studies described above revealed several benefits of applying AI for lactate monitoring even though the latter is not without drawbacks. One of the main issues is the ability to incorporate AI models into practices of intensive care medicine and EHR platforms. Some ICUs may not have the necessary set up of processing real-time data and integrating Artificial Intelligence in their affair which may lessen the capacity of such models. Furthermore, artificial intelligence models are often deployed after a clinician undertakes the process of interpreting predictions and incorporating the AI findings into the overall decision-making [42]. Another one is the difficulty in determining interpretability and transparency of the data on which AI makes its predictions. While models like decision trees tell how variables affect some predictions, other complex models like neural networks are hard to explain. If clinicians are to rely and put their trust on AI-generated alerts, there must be explain ability or interpretability of the model's decision making. Solving this problem entails adopting dual-stage frameworks where researchers use easy-to-understand models that elucidate results alongside tough-to-interpret models that produce accurate predictions. Finally, ethical and regulatory issues are important in relation to the patients' information privacy and security. AI models depend on the patient's data, therefore, patient data must meet certain guidelines such as the HIPAA in the United States [43]. Hospitals and those providing technologies require that these form strict measures of compliance thus having good policies for the privilege of data all for the sake of protecting patients' information.

This paper discusses the clinical uses and specific cases of applying AI in lactate monitoring, which demonstrate the effectiveness of creating predictive models in enhancing the treatment required in critical care. Real-time data in the form of predictive analysis gives clinicians a proactive role, as errors require corrections from patients' timelines, thereby giving patients shorter recovery periods [44]. While there are some issues regarding implementation, data integration and other important regulatory questions that have not yet found a satisfactory solution, the success gained in various case studies show the importance of AI as a tool for improving the quality of intensive care. In the longer term, we anticipate that growth in both AI technology and healthcare system will lead to the more common use of AI-supported lactate monitoring and the establishment of novel benchmarks for patient safety and more efficient clinical care for critical care patients [45].

HYPOTHETICAL DEVELOPMENTS, CAUTIONS, AND POTENTIAL RISK LESSONS ON ARTIFICIAL INTELLIGENCE IN LACTATE MONITORING

Looking at the future directions and challenges of AI and ML in the healthcare especially in critical care monitoring the following are evident. Artificial intelligence based lactate monitoring, shown to have value in influencing patient improvement in the aftermath of cardiac surgery by enabling earlier intervention and individualization of treatment. However, when these systems are being transposed from research prototypes to real-world clinical applications, a range of issues must be taken into account to guarantee the proper, safe, and rights compliant use [46]. In this section, the authors outline the prospective lines of development in the further work on lactate monitoring, describe the problems of AI implementation and regulation related to lactate modeling and share the considerations on the ethical consequences of using such sophisticated technologies in the health care field.

Directions for future studies in monitoring lactate using AI: The current state of research for using AI in lactate monitoring shows that development for this application is highly likely to move quickly as improvements are made on the technologies being used and the different algorithms, data possibilities, and patient-based uses are explored. A new paradigm is to proposed use of Comprehensive AI Models of Patients outside the standard Intensive Care Unit parameters [47]. For example, incorporating genomics, real time imaging and environmental data could result in more accurate and comprehensive models that could now predict not only lactate levels but also many other potential post-surgical complications. Next generation data will complement different data types, while future AI models could help provide more informative assessment of patient health, moving CARE toward more precision medicine approaches [48].

A second area for future research includes work that will address issues concerning constant model updating and model flexibility. The current collection of AI models is mainly trained on the past and fixed once in use much of the time. Thus, to keep the accuracy constant with time and when data is collected from variable intensive care unit environments, there is a need for models that are capable of adjusting with time as more patient data is processed and new trends are characterized. Further studies of reinforcement learning a sub type of machine learning in operation with feedback information that is continually updated, would allow for models that adapt to changes in patient conditions and therefore more flexibility in exceptional monitoring [49]. This raises a number of questions around the technical and operational feasibility of implementing Lactate/AI to scale, in reality. One of the key challenges is the ability of the AI leadership models in integrating with hospital settings, as well as integrating the EHR systems and ICU equipment. To achieve perpetuity in the way data is fed to these models, many hospitals and clinics use the various platforms; making it tough to standardize data. To reliably transmit and process data emanating from various monitoring instruments in real-time for analyses that enable prediction of an impending situation poses certain challenges in terms of capital investment and staff training [50].





One more technical issue arises when it is required to develop sound AI systems excluding adverse health effects for a wide range of patients. It is also important to note that AI models can have institutional bias and this is mainly due to the limited data with which the AI models are trained from some diverse demographic or clinic type. Being able to ensure that these models can easily generalize across different populations which are equally not biased remains an important factor. Because of this, rigorous testing, and model validation in other healthcare facilities are critical again before widespread implementation [51]. Additionally, the parameter of “alarm fatigue” is an issue in this case. There is also the risk that in an ICU context and especially when an AI model has been deployed to notify clinicians of lactate increase or any other complication, the alert fatigue may set in and clinicians receive alerts they deemed irrelevant. If this is not addressed, alarm fatigue results in reduced response to important alerts to be given to the clinician. The fine tuning of such systems to reduce loud bells that do not add value to the care of the patient without providing a negative impact on patient care is an important feature to consider when implementing artificial intelligence.


AI in the health sector is lightly regulated based in existing health laws, but faces several ethical issues including data sharing and reporting. One of the major ethical considerations is to secure the identity of their patients while at the same time dealing with a massive amount of health information [52]. Applications of AI models in lactate tracking depend on the frequent data feed such as vitals and labs signs that are considered as sensitive data under the Hipaa regulations. There is also a paramount need that ensure data is collected, stored and processed in a secure manner to answer legal requirements and patient’s trust. Another problem of ethical concern is transparency. The problem is, many current advanced AI models including deep learning algorithms are often even described as ‘black boxes’ because one cannot interpret the algorithmic decisions. This lack of transparency can be an issue in healthcare as clinicians need to be able to understand the thought process behind AI created prediction in order to make appropriate decisions. To overcome this there is research into solutions including the integration of transparent models with deep learning allowing clinicians to understand how features have informed predictions made by the model [53]. Two other ethical issues are accountability and liability. In the case of an AI model where inaccurate predictions result in negative consequences identifying the degree of fault is not easy. Trying to establish AI’s role as a decision support tool and not an independent decision-maker is important to define boundaries for clinicians and other health care professionals on their use of AI driven outputs.

Accessibility and Non-Bias: With advances in and implementation of AI-based lactate monitoring, issues concerning fairness of access of such systems and opportunities for bias in these models exist. This is an issue of health inequality; the application of AI and especially the latest advanced technology can be challenging in hospitals which cannot afford or lack the infrastructure for implementing AI solutions [54]. If AI-driven monitoring tools are only available or made available first in high resource settings, this could only elongate existing healthcare disparities. Therefore, there is a need to design AI solutions that are cheap and can fit any healthcare system regardless of its healthcare technological sophistication. Other is that AI models themselves can be biased and thus can lead to inequities in patients’ outcomes. Should AI models be trained in specific populations and not diverse, the models are likely to provide less accurate results to the minority’s population. Ensuring no bias is easier said than done it involves using diverse training datasets, undergoing several validations and even retesting models to help in ensuring that the AI models offer appropriate predictions for all classes of people.

CONCLUSION

The ability to use an information system with artificial intelligence to predict and managing blood lactic acid levels following cardiac surgery is an innovative area in critical care that provides a proactive approach to patient care. This technology changes the paradigm from fixed monitoring where patient’s condition is checked at intervals to Monitor-Check-Act kind of monitoring where care is provided as soon as an abnormality in the patient’s condition is detected likely reducing the complications caused by high levels of lactate. Through the use of large quantities of physiological data, an AI model is able to identify faint signs of the patient’s metabolic stress and enable the clinician to intervene before the situation escalates and the patient is admitted into the ICU. Innovations in AI and machine learning are opening up possibilities of developing complex, combinational models to enable integration of more elaborate data formats in order to improve the precision of the models. In this paper, a number of real-life examples prove the efficacy of using artificial intelligence concerning lactate levels to predict the development of complications, tailor treatment, and enhance ICU efficiency.

However, some barriers remain, get-toing with the infrastructure of a hospital, a lack of standardization and that problem of alarm fatigue again. Moreover, there are important questions of ethical nature including but not limited to privacy, unpredictability, and liability concerning patients. There is a need to ensure that these systems are explainable and effective in developing trust and that no unfair inequalities will be created between groups of the society as regards to patients. Considering the future advancement of AI-based lactate monitoring, critical care has the best chance at improving and moving forward. Nevertheless, as a result of further investigations, cooperation with other disciplines, and compliance with ethical standards, these instruments can become the basis for the continued development of individualized medicine in the field of postoperative rehabilitation. To unlock the full benefit of AI, resolving the technical, operational, and moral issues will be the moving force to the next level. As healthcare systems shift to the use of artificial intelligence solutions for disease diagnostics and treatment, normal lactate monitoring as demonstrated here can develop a new culture of






patient-centered, efficient, and effective personalized medicine to improve the quality and equity of ICU care across culturally diverse populations.


REFERENCES

1. Hoste EAJ, Kellum JA, Selby NM et al (2018) Global epidemiology and outcomes of acute kidney injury. *Nat Rev Nephrol* 14(10):607–625. <https://doi.org/10.1038/s41581-018-0052-02>.
2. Van den Eynde J, Delpire B, Jacquemyn X et al (2022) Risk factors for acute kidney injury after pediatric cardiac surgery: a meta-analysis. *Pediatr Nephrol* 37(3):509–519. <https://doi.org/10.1007/s00467-021-05297-0>
3. Webb TN, Goldstein SL (2017) congenital heart surgery and acute kidney injury. *Curr Opin Anaesthesiol* 30(1):105–112. <https://doi.org/10.1097/ACO.0000000000000406>
4. Ricci Z, Raggi V, Marinari E et al (2022) acute kidney injury in pediatric cardiac intensive care children: not all admissions are equal: a retrospective study. *J Cardiothorac Vasc Anesth* 36(3):699–706. <https://doi.org/10.1053/j.jvca.2021.04.015>
5. Romagnoli S, Ricci Z (2015) Postoperative acute kidney injury. *Minerva Anestesiol* 81(6):684–696
6. Meersch M, Schmidt C, Hofmeier A et al (2017) (2017) Prevention of cardiac surgery-associated AKI by implementing the KDIGO guidelines in high risk patient's identified by biomarkers: the PrevAKI randomized controlled trial [published correction appears in *Intensive Care Med*. *Intensive Care Med* 43(11):1551–1561. <https://doi.org/10.1007/s00134-016-4670-3>
7. Bellos I, Iliopoulos DC, Perrea DN (2019) Pharmacological interventions for the prevention of acute kidney injury after pediatric cardiac surgery: a network meta-analysis. *Clin Exp Nephrol* 23(6):782–791. <https://doi.org/10.1007/s10157-019-01706-9>
8. Kelly BJ, Chevarria J, O'Sullivan B, Shorten G (2021) the potential for artificial intelligence to predict clinical outcomes in patients who have acquired acute kidney injury during the perioperative period. *Perioper Med (Lond)* 10(1):49. <https://doi.org/10.1186/s13741-021-00219-y>
9. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative (2008) The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 61(4):344–9. <https://doi.org/10.1016/j.jclinepi.2007.11.008>
10. Neugarten J, Sandilya S, Singh B, Golestaneh L. Sex and the risk of AKI following cardio-thoracic surgery: a meta-analysis. *Clin J Am Soc Nephrol*. 2016; 11:2113–22.
11. James MT, Bhatt M, Pannu N, Tonelli M. Long-term outcomes of acute kidney injury and strategies for improved care. *Nat Rev Nephrol*. 2020; 16:193–205.
12. Pickkers P, Darmon M, Hoste E, Joannidis M, Legrand M, Ostermann M, et al. Acute kidney injury in the critically ill: an updated review on pathophysiology and management. *Intensive Care Med*. 2021; 47:835–50.
13. Schunk SJ, Zarbock A, Meersch M, Küllmar M, Kellum JA, Schmit D, et al. Association between urinary dickkopf-3, acute kidney injury, and subsequent loss of kidney function in patients undergoing cardiac surgery: an observational cohort study. *Lancet*. 2019; 394:488–96
14. Ugwuowo U, Yamamoto Y, Arora T, Saran I, Partridge C, Biswas A, et al. Real-time prediction of acute kidney injury in hospitalized adults: implementation and proof of concept. *Am J Kidney Dis*. 2020; 76:806–14?
15. Wijesundera DN, Karkouti K, Dupuis JY, Rao V, Chan CT, Granton JT, et al. Derivation and validation of a simplified predictive index for renal replacement therapy after cardiac surgery. *JAMA*. 2007; 297:1801–9.
16. Mehta RH, Grab JD, O'Brien SM, Bridges CR, Gammie JS, Haan CK, et al. Bedside tool for predicting the risk of postoperative dialysis in patients undergoing cardiac surgery. *Circulation*. 2006; 114:2208–16.
17. Jiang W, Xu J, Shen B, Wang C, Teng J, Ding X. Validation of four prediction scores for cardiac surgery-associated acute kidney injury in chinese patients. *Braz J Cardiovasc Surg*. 2017; 32:481–6.
18. Tomašev N, Glorot X, Rae JW, Zielinski M, Askham H, Saraiva A, et al. A clinically applicable approach to continuous prediction of future acute kidney injury. *Nature*. 2019; 572:116–9.
19. Lundberg SM, Erion GG, Lee S-I. Consistent individualized feature attribution for tree ensembles. *arXiv [Preprint]*. 2018. <https://arxiv.org/abs/1802.03888>. Accessed 12 Feb 2018.
20. Collins GS, Reitsma JB, Altman DG, Moons KG. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD statement. *BMJ*. 2015; 350:g7594.
21. Stevens PE, Levin A. Evaluation and management of chronic kidney disease: synopsis of the kidney disease: improving global outcomes 2012 clinical practice guideline. *Ann Intern Med*. 2013; 158:825–30.
22. Chen T, Guestrin C. XGBoost: A Scalable Tree Boosting System. *arXiv [Preprint]*. 2016. <https://arxiv.org/abs/1603.02754>. Accessed 9 Mar 2016





23. Zhou ZH, Feng J. Deep Forest: towards an alternative to deep neural networks. arXiv [Preprint]. 2017. <https://arxiv.org/abs/1702.08835v1>. Accessed 28 Feb 2017
24. Niculescu-Mizil A, Caruana R. Predicting good probabilities with supervised learning. In: Machine learning: Proceedings of the 22nd international conference. 2005. See section 4 (Qualitative Analysis of Predictions).
25. Shapley LS. A value for n-person games. Princeton: Princeton University Press; 1953. 19. Nagin DS, Jones BL, Passos VL, Tremblay RE. Group-based multi-trajectory modeling. *Stat Methods Med Res.* 2018; 27:2015–23.
26. Cook NR. Statistical evaluation of prognostic versus diagnostic models: beyond the ROC curve. *Clin Chem.* 2008; 54:17–23.
27. Vickers AJ, Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. *Med Decis Making.* 2006; 26:565–74.
28. Kanji HD, Schulze CJ, Hervas-Malo M, Wang P, Ross DB, Zibdawi M, et al. Difference between pre-operative and cardiopulmonary bypass mean arterial pressure is independently associated with early cardiac surgery-associated acute kidney injury. *J Cardiothorac Surg.* 2010; 5:71.
29. O'Neal JB, Shaw AD, Billings FT. Acute kidney injury following cardiac surgery: current understanding and future directions. *Crit Care.* 2016; 20:187.
30. Gambardella I, Gaudino M, Ronco C, Lau C, Ivascu N, Girardi LN. Congestive kidney failure in cardiac surgery: the relationship between central venous pressure and acute kidney injury. *Interact Cardiovasc Thorac Surg.* 2016; 23:800–5.
31. Lopez MG, Shotwell MS, Morse J, Liang Y, Wanderer JP, Absi TS, et al. Intraoperative venous congestion and acute kidney injury in cardiac surgery: an observational cohort study. *Br J Anaesth.* 2021; 126:599–607.
32. Manning MW, Li YJ, Linder D, Haney JC, Wu YH, Podgoreanu MV, et al. Conventional ultrafiltration during elective cardiac surgery and postoperative acute kidney injury. *J Cardiothorac Vasc Anesth.* 2021; 35:1310–8.
33. Tseng PY, Chen YT, Wang CH, Chiu KM, Peng YS, Hsu SP, et al. Prediction of the development of acute kidney injury following cardiac surgery by machine learning. *Crit Care.* 2020; 24:478.
34. Esteve F, Lopez-Delgado JC, Javierre C, Skaltsa K, Carrio ML, RodríguezCastro D, et al. Evaluation of the PaO₂/FiO₂ ratio after cardiac surgery as a predictor of outcome during hospital stay. *BMC Anesthesiol.* 2014; 14:83.
35. Gao XP, Zheng CF, Liao MQ, He H, Liu YH, Jing CX, et al. Admission serum sodium and potassium levels predict survival among critically ill patients with acute kidney injury: a cohort study. *BMC Nephrol.* 2019; 20:311
36. Zhang D, Shi L. Serum lactate dehydrogenase level is associated with inhospital mortality in critically ill patients with acute kidney injury. *Int Urol Nephrol.* 2021; 53:2341–8.
37. Kym D, Cho YS, Yoon J, Yim H, Yang HT. Evaluation of diagnostic biomarkers for acute kidney injury in major burn patients. *Ann Surg Treat Res.* 2015; 88:281–8.
38. Guan C, Li C, Xu L, Zhen L, Zhang Y, Zhao L, et al. Risk factors of cardiac surgery-associated acute kidney injury: development and validation of a perioperative predictive nomogram. *J Nephrol.* 2019; 32:937–45.
39. Vermeulen Windsant IC, Snoeijs MG, Hanssen SJ, Altintas S, Heijmans JH, Koepfel TA, et al. Hemolysis is associated with acute kidney injury during major aortic surgery. *Kidney Int.* 2010; 77:913–20
40. Weedle RC, Da Costa M, Veerasingam D, Soo AWS. The use of neutrophil lymphocyte ratio to predict complications post cardiac surgery. *Ann Transl Med.* 2019; 7:778.
41. Wang Q, Li J, Wang X. The neutrophil-lymphocyte ratio is associated with postoperative mortality of cardiac surgery. *J Thorac Dis.* 2021; 13:67–75
42. Haran C, Gimpel D, Clark H, McCormack DJ. Preoperative neutrophil and lymphocyte ratio as a predictor of mortality and morbidity after cardiac surgery. *Heart Lung Circ.* 2021; 30:414–8.
43. Smith LE, Smith DK, Blume JD, Linton MF, Billings FT 4th. High-density lipoprotein cholesterol concentration and acute kidney injury after cardiac surgery. *J Am Heart Assoc.* 2017; 6:e006975.
44. Rhodes JF, Blaufox AD, Seiden HS, Asnes JD, Gross RP, Rossi AF (1999) Cardiac arrest in infants after congenital heart surgery. *Circulation* 100(19 Suppl): II 194-9. https://doi.org/10.1161/01.cir.100.suppl_2.ii-194. (PMID: 10567303)
45. Ichord R, Silverstein FS, Slomine BS, Telford R, Christensen J, Holubkov R, Dean JM, Moler FW (2018) THAPCA Trial Group Neurologic outcomes in pediatric cardiac arrest survivors enrolled in the THAPCA trials. *Neurology* 91(2):e123–e131. <https://doi.org/10.1212/WNL.0000000000005773>
46. Alten JA, Klugman D, Raymond TT et al (2017) Epidemiology and outcomes of cardiac arrest in pediatric cardiac ICUs. *Pediatr Crit Care Med* 18(10):935–943





47. Husnain, A., Alomari, G., & Saeed, A. AI-Driven Integrated Hardware and Software Solution for EEG-Based Detection of Depression and Anxiety.
48. Alten J, Cooper DS, Klugman D et al (2022) Preventing cardiac arrest in the pediatric cardiac intensive care unit through multicenter collaboration. *JAMA Pediatr* 176(10):1027–1036. <https://doi.org/10.1001/jamapediatrics.2022.2238>
49. Ferguson LP, Thiru Y, Stafa SJ et al (2020) Reducing cardiac arrests in the PICU: initiative to improve time to administration of prearrest bolus epinephrine in patients with cardiac disease. *Crit Care Med* 48(7):e542–e549. <https://doi.org/10.1097/ccm.0000000000004349>
50. Riley CM, Murphy LD, Mastropietro CW (2022) Cardiac arrest in children following cardiac surgery: a scoping review of contributing factors. *World J Pediatr Congenit Heart Surg* 13(4):475–481. <https://doi.org/10.1177/21501351221100791>
51. Khoubian JJ, Alten JA (2022) Cardiac arrest in pediatric cardiac ICUs: prevention comes first. *Curr Treat Options Peds* 8:325–333. <https://doi.org/10.1007/s40746-022-00255-w>
52. Alten J, Cooper DS, Klugman D et al (2022) Preventing cardiac arrest in the pediatric cardiac intensive care unit through multicenter collaboration. *JAMA Pediatr* 176(10):1027–1036. <https://doi.org/10.1001/jamapediatrics.2022.2238>
53. Shin Y, Cho KJ, and Lee Y et al (2022) Multicenter validation of a deep-learning-based pediatric early warning system for prediction of deterioration events. *Acute Crit Care* 37(4):654–666. <https://doi.org/10.4266/acc.2022.00976>
54. Husnain, A., & Saeed, A. (2024). AI-enhanced depression detection and therapy: Analyzing the VPSYC system. *IRE Journals*, 8(2), 162-168.

